

On bilinear algorithms for multiplication in quaternion algebras

Vladimir Lysikov*
Moscow State University

Abstract

We show that the bilinear complexity of multiplication in a non-split quaternion algebra over a field of characteristic distinct from 2 is 8. This question is motivated by the problem of characterising algebras of almost minimal rank studied in [1]

This paper is a translation of a report submitted by the author to the XI international seminar "Discrete mathematics and its applications".

Definition 1. A sequence $(f_1, g_1, z_1; \dots; f_r, g_r, z_r)$ with $f_k \in U^*, g_k \in V^*, z_k \in W$ is called a *bilinear algorithm* of length r for a bilinear mapping $\varphi: U \times V \rightarrow W$ if

$$\varphi(u, v) = \sum_{k=1}^r f_k(u)g_k(v)z_k \quad \forall u \in U, v \in V. \quad (1)$$

The minimal length of a bilinear algorithm for φ is called the *bilinear complexity*, or the *rank*, of φ and is denoted by $R(\varphi)$. Algorithms of minimal length are called optimal.

Rank of bilinear map has a useful algebraic interpretation. A bilinear map can be thought of as a tensor in $U^* \otimes V^* \otimes W$. Bilinear algorithms are decompositions of this tensor and the bilinear complexity coincides with the tensor rank as defined in multilinear algebra.

Definition 2. A quaternion algebra over field F of characteristic $\text{char } F \neq 2$ is a 4-dimensional algebra generated by two elements i, j with identities

$$i^2 = p, \quad j^2 = q, \quad ij = -ji \quad (2)$$

for some $p, q \in F^\times$.

Any quaternion algebra over F is either isomorphic to the matrix algebra $F^{2 \times 2}$ (in this case it is called split) or a noncommutative division algebra. It is known that $R(F^{2 \times 2}) = 7$ and $R(H) \geq 8$ for non-split quaternion algebra H [3].

Let $m = \dim U, n = \dim V$. We will prove some general results on algorithms of rank $m + n$ for some class of bilinear maps.

Definition 3. We say that an element $u_0 \in U$ is (left) φ -regular if the linear map $\varphi(u_0, \cdot)$ is an injection, i. e., if

$$\varphi(u_0, v) = 0 \Leftrightarrow v = 0. \quad (3)$$

Definition 4. A bilinear algorithm $\varphi = \sum_{k=1}^r f_k \otimes g_k \otimes z_k$ is called *two-component* if the set $\{1, \dots, r\}$ of indices can be partitioned into two sets I and J such that $\{f_i | i \in I\}$ and $\{g_j | j \in J\}$ are bases of U^* and V^* respectively.

Lemma 1. If $R(\varphi) = m + n$, $\text{lker } \varphi = \mathbf{0}$ and every basis of U contains a φ -regular element then all optimal bilinear algorithms for φ are two-component.

Proof. Let $\sum_{k=1}^{m+n} f_k \otimes g_k \otimes z_k$ be an optimal bilinear algorithm for φ . Since $\text{lker } \varphi = \mathbf{0}$, the functionals $f_1, \dots, f_r$ span U^* . W.l.o.g. let $f_1, \dots, f_m$ be a basis and u_1 be a φ -regular element of the dual basis $u_1, \dots, u_n$. Since u_1 is regular, functionals $g_1, g_{m+1}, \dots, g_{m+n}$ span V^* .

Case 1. $g_{m+1}, \dots, g_{m+n}$ are linearly independent. Then $I = \{1, \dots, m\}$ and $J = \{m+1, \dots, m+n\}$ form a partition of $\{1, \dots, m+n\}$ required by definition of two-component algorithm.

Case 2. $g_{m+1}, \dots, g_{m+n}$ are linearly dependent. In this case $\dim \text{lin}(\{g_{m+1}, \dots, g_{m+n}\}) = n - 1$ and there is a unique (up to a constant) linear dependence of $g_{m+1}, \dots, g_{m+n}$. W.l.o.g. assume that $\sum_{k=1}^s c_k g_{m+k} = 0$ where all the coefficients c_k are nonzero.

*E-mail: lysikov-vv@yandex.ru

If there is an index p such that $m+1 \leq p \leq m+s$ and f_p has a nonzero first coordinate in basis $f_1, \dots, f_m$ then $I = \{2, \dots, m, p\}$ and $J = \{1, m+1, \dots, m+n\} \setminus \{p\}$ form a required partition. If all $f_{m+1}, \dots, f_{m+s}$ have zero first coordinates, then we have a contradiction with φ -regularity of u_1 , since $\varphi(u_1, v) = 0$ for a nonzero $v \in \ker g_1 \cap \bigcap_{k=n+s+1}^{m+n} \ker g_k$. $\square$

Lemma 2. *A two-component bilinear algorithm for φ exists if and only if there are bases $(u_1, \dots, u_m)$ and $(v_1, \dots, v_m)$ of spaces U and V resp. and collections $(z'_1, \dots, z'_m)$, $(z''_1, \dots, z''_n)$ with $z'_i, z''_j \in W$ such that the following condition holds:*

$$\varphi(u_i, v_j) \in \text{lin}(\{z'_i, z''_j\}). \quad (4)$$

Proof. Any two-component bilinear algorithm for φ can be written as

$$\varphi = \sum_{i=1}^m f_i \otimes \left(\sum_{j=1}^n \lambda_{ij} g_j \right) \otimes z'_i + \sum_{j=1}^n \left(\sum_{i=1}^m \mu_{ij} f_i \right) \otimes g_j \otimes z''_j, \quad (5)$$

where (f_i) and (g_j) are bases of U^* and V^* respectively. Let (u_i) and (v_j) be the bases dual to (f_i) and (g_j) . Then it follows that

$$\varphi(u_i, v_j) = \lambda_{ij} z'_i + \mu_{ij} z''_j. \quad (6)$$

Conversely, if (6) holds then (5) is a two-component algorithm for φ . $\square$

Corollary. *Let A be a local algebra, $\dim A = n$, and $R(A) \geq 2n$. Equality $R(A) = 2n$ holds iff there are bases $(u_1 = 1, u_2, \dots, u_n)$, $(v_1 = 1, v_2, \dots, v_n)$ and collections $(z'_1, \dots, z'_n)$, $(z''_1, \dots, z''_n)$ such that*

$$u_i v_j \in \text{lin}(\{z'_i, z''_j\}). \quad (7)$$

Theorem 1. *Let F be a field of characteristic $\text{char } F \neq 2$ and H be a non-split quaternion algebra over F . Then $R(H) = 8$.*

Proof. Apply the preceding corollary with the following bases and collections:

$$(u_1, u_2, u_3, u_4) = (v_1, v_2, v_3, v_4) = (1, i, j, k) \quad (8a)$$

$$(z'_1, z'_2, z'_3, z'_4) = (1 + \alpha i + \beta j + \gamma k, 1 + \alpha i - \beta j - \gamma k, 1 - \alpha i + \beta j - \gamma k, 1 - \alpha i - \beta j + \gamma k) \quad (8b)$$

$$(z''_1, z''_2, z''_3, z''_4) = (1 - \alpha i - \beta j - \gamma k, 1 - \alpha i + \beta j + \gamma k, 1 + \alpha i - \beta j + \gamma k, 1 + \alpha i + \beta j - \gamma k) \quad (8c)$$

where $\alpha, \beta, \gamma \in F$, $\alpha \neq 0$, $\beta \neq 0$, $\gamma \neq 0$. $\square$

By considering bilinear algorithms arising from different constants α, β, γ we also managed to prove the following fact about equivalence of optimal algorithms in the sense of de Groote [2].

Theorem 2. *Let F be a field of characteristic $\text{char } F \neq 2$ and H be a non-split quaternion algebra over F . There are infinitely many de Groote equivalence classes of optimal bilinear algorithms for the multiplication in H .*

The author is grateful to Prof. V. B. Alekseyev for his attention to this research.

This research was supported by RFBR grant 12-01-91331-DFG-a.

Bibliography

1. M. Bläser, A. M. de Voltaire. Semisimple algebras of almost minimal rank over the reals // Theoretical Computer Science. — 2009. — vol. 410, no. 50. — pp. 5202–5214.
2. H. F. de Groote. On varieties of optimal algorithms for the computation of bilinear mappings I // Theoretical Computer Science. — vol. 7, no. 1. — pp. 1–24.
3. P. Bürgisser, M. Clausen, M. A. Shokrollahi. Algebraic complexity theory. — Springer Verlag, 1997.